

Spontaneous generation of a half-quantum vortex in spin-orbit coupled Bose-Einstein condensates

Congjun Wu¹ and Ian Mondragon-Shem^{1,2}

¹*Department of Physics, University of California, San Diego, CA 92093*

²*Instituto de Física, Universidad de Antioquia, AA 1226, Medellín, Colombia*

According to the “*no-node*” theorem, many-body ground state wavefunctions of conventional Bose-Einstein condensation (BEC) are positive-definite, thus time-reversal symmetry cannot be spontaneously broken. We find that multi-component bosons with spin-orbit coupling provide an unconventional type of BEC beyond this paradigm. In a harmonic confining trap, the condensate spontaneously generates a half-quantum vortex combined with the skyrmion type of spin texture. This phenomenon can be realized for both the cold atom systems with artificial spin-orbit coupling generated from atom-laser interaction, and for exciton condensates in semi-conductor systems.

PACS numbers: 71.35.-y, 73.50.-h, 03.75.Mn, 03.75.Nt

The conventional many-body ground state wavefunctions of bosons satisfy the celebrated “*no-node*” theorem in the absence of rotation, as pointed out by Feynman [1], which means that they are positive-definite in the coordinate representation. This theorem implies that time-reversal (TR) symmetry cannot be spontaneously broken. It applies to various quantum ground states of bosons including Bose-Einstein condensates (BEC), Mott-insulating states, density-wave states, and super-solid states, thus making this a very general statement.

It would be interesting to search for novel types of quantum states of bosons beyond this paradigm, such as unconventional BECs with complex-valued wavefunctions. The “*no-node*” theorem is only a ground state property, hence it does not apply to excited states. Recently, a meta-stable state of bosons has been realized, using cold atoms in optical lattices, by pumping atoms into high orbital bands (e.g. *p*-bands) with a long life time [2, 3]. Interactions among orbital bosons are characterized by an orbital Hund’s rule [4, 5], which gives rise to a class of complex superfluid states by developing onsite orbital angular momentum, thus breaking TR symmetry spontaneously [3–7].

The “*no-node*” theorem does not apply to spinful bosons with spin-orbit (SO) coupling either. The linear dependence on momentum introduced by the SO coupling invalidates Feynman’s proof. Although alkali atoms are too heavy to exhibit relativistic SO coupling, artificial SO coupling can be induced by laser-atom interactions [8–13]. The BECs in such systems have been studied, and have been shown to exhibit frustrations of superfluidity [14]. On the other hand, excitons in semiconductors [15–18] with small effective mass can exhibit SO coupling in the center-of-mass motion [19–21]. In particular, exciting progress has been made in indirect exciton systems in coupled quantum wells where electrons and holes are spatially separate [22–25]. The extraordinarily long life time of indirect excitons provides a wonderful opportunity to investigate the exotic state of matter of exciton condensation [16].

In this article, we show that bosons with Rashba SO coupling provide a new opportunity to investigate unconventional BEC beyond the “*no-node*” theorem. We find that Rashba SO coupling leads to fragmented condensation in homogeneous systems. Furthermore, with an external harmonic potential trap, TR symmetry is spontaneously broken. In this case, the ground state condensate develops orbital angular momentum, which can be viewed as a half-quantum vortex. Moreover, the spin density distribution exhibits a cylindrically symmetric spiral pattern as skyrmions. The experimental realization, in both cold atom and semiconductor exciton systems, is discussed.

We begin with a 3D two-component boson system with Rashba SO coupling in the *xy*-plane and with contact interaction, described by

$$H = \int d^3\vec{r} \left[\psi_\alpha^\dagger \left[-\frac{\hbar^2 \nabla^2}{2M} - \mu \right] \psi_\alpha + \hbar \lambda_R \psi_\alpha^\dagger [-i \nabla_y \sigma_x + i \nabla_x \sigma_y] \psi_\beta + \frac{g}{2} \psi_\alpha^\dagger \psi_\beta^\dagger \psi_\beta \psi_\alpha \right], \quad (1)$$

where ψ_α is the boson operator; the pseudospin indices $\alpha = \uparrow, \downarrow$ refer to two different internal components of the bosons; λ_R is the SO coupling strength; g describes the *s*-wave scattering interaction. Although Eq. (1) is for bosons, it satisfies a suitably defined TR symmetry $T = i\sigma_2 C$ satisfying $T^2 = -1$, where C is the complex conjugate operation and σ_2 operates on the boson pseudospin degree of freedom.

In the homogeneous system, the single particle states are the helicity eigenstates of $\vec{\sigma} \cdot (\vec{k} \times \hat{z})$ which have a dispersion relation given by $\epsilon_\pm(\vec{k}) = \frac{\hbar^2}{2M} ((k - k_0)^2 + k_z^2)$, where $k_0 = \frac{M\lambda_R}{\hbar}$. The energy minima are located on the lower branch along a ring with radius k_0 in the plane of $k_z = 0$. The corresponding two-component wavefunction $\psi_+(\vec{k})$ with $|\vec{k}| = k_0$ can be solved as $\psi_+^T(\vec{k}) = \frac{1}{\sqrt{2}}(e^{-i\phi_k/2}, ie^{i\phi_k/2})$, where ϕ_k is the azimuthal angle of the projection of $\vec{k}$ in the *xy*-plane. The interaction part in Eq. (1) in the helicity basis can be




FIG. 1: The degeneracy of the single particle ground states along a circle with the radius k_0 in momentum space. We assume that the fragmented condensation occurs at points A ($k_0\hat{e}_x, 0, 0$) and B ($-k_0\hat{e}_x, 0, 0$) with spin polarizations along $\pm\hat{e}_y$, respectively. The low energy excitations within $|k-k_0| < \Lambda$, $|k_z| < \Lambda$, and $\Lambda/k_0 \ll 1$ are classified into two regimes I (inside two cylinders centering around A and B with radius of Λ) and II (outside).

represented as

$$H_{int} = \frac{g}{2} \sum_{\lambda\mu\nu\rho} \sum_{p_1 p_2 q} \langle \vec{p}_1 + \vec{q}; \lambda | \vec{p}_1; \rho \rangle \langle \vec{p}_2 - \vec{q}; \mu | \vec{p}_2; \nu \rangle \times \psi_\lambda^\dagger(\vec{p}_1 + \vec{q}) \psi_\mu^\dagger(\vec{p}_2 - \vec{q}) \psi_\nu(\vec{p}_2) \psi_\rho(\vec{p}_1), \quad (2)$$

where the Greek indices λ, ν, μ, ρ are the helicity indices $\pm$; the matrix elements denote the inner product of spin wavefunctions of two helicity eigenstates at different momenta, e.g., $\langle \vec{p}_1 + \vec{q}; \lambda | \vec{p}_1; \rho \rangle = \frac{1}{2}[1 + \lambda\rho e^{i(\phi_{p_1} - \phi_{p_1+q})}]$.

At the Hartree-Fock level, the kinetic energy can be minimized through coherent or fragmented condensation in the linear superpositions of the single particle ground states. For the interaction energy, the exchange part among bosons is positive within the s -wave scattering approximation where g is independent of momentum, thus the ground state favors the zero exchange energy. The spin polarizations at $\vec{k}$ and $-\vec{k}$ in the plane of $k_z = 0$ are orthogonal to each other. Thus, without loss of generality, the fragmented condensate defined below satisfies this requirement

$$\Phi_0 = \frac{1}{\sqrt{N_A! N_B!}} [\psi_+^\dagger(\vec{k}_A)]^{N_A} [\psi_+^\dagger(\vec{k}_B)]^{N_B} |0\rangle, \quad (3)$$

where $\vec{k}_A = (-k_0, 0, 0)$ and $\vec{k}_B = (k_0, 0, 0)$; and (N_A, N_B) is the particle number partition satisfying $N_A + N_B = N_0$, with N_0 the total particle number in the condensate. In homogeneous systems, fragmented condensates with different partitions do not mix because of momentum conservation.

An “order from disorder” calculation on the low energy excitation spectra is needed to determine the optimal particle number partition. The momentum cutoff $\hbar\Lambda$ is defined as $\hbar^2\Lambda^2/(2M) \approx gn$, where gn is the interaction energy scale and n is the particle density. We only consider the limit of strong SO coupling, $k_0 \gg \Lambda$, and leave

the general case for later study. The low energy Bogoliubov excitations that significantly mix particle and hole operators mainly lie in the regime $|\sqrt{k_x^2 + k_y^2} - k_0| < \Lambda$ and $|k_z| < \Lambda$, which can be further divided into two parts I and II as depicted in Fig.1. Part I is inside two cylinders with the radius of Λ centering around A and B, and part II is outside these two cylinders.

For the low energy states in part I, we define the bosonic operators in the lower branch as $a_{\vec{q}} = \psi_+(-k_0\hat{e}_x + \vec{q})$ and $b_{\vec{q}} = \psi_+(k_0\hat{e}_x + \vec{q})$, and $\langle a_{q=0} \rangle = \sqrt{N_A}$ and $\langle b_{q=0} \rangle = \sqrt{N_B}$ respectively. The relative phase between two condensates can be set to zero due to a suitable translation operation. The low energy excitations in this region have been calculated in Ref. [14]. By defining $\gamma_1^\dagger(\vec{q}) = \frac{1}{N_0}(\sqrt{N_A}a_{\vec{q}}^\dagger + \sqrt{N_B}b_{\vec{q}}^\dagger)$, $\gamma_2^\dagger(\vec{q}) = \frac{1}{N_0}(\sqrt{N_B}a_{\vec{q}}^\dagger - \sqrt{N_A}b_{\vec{q}}^\dagger)$, the mean-field Hamiltonian in region I, up to the quadratic level of q , can be represented as

$$H_{MF,1} = \sum_{\vec{q}} E(\vec{q}) \{ \gamma_1^\dagger(\vec{q}) \gamma_1(\vec{q}) + \gamma_2^\dagger(\vec{q}) \gamma_2(\vec{q}) \} + gN_0 \{ \gamma_1^\dagger(\vec{q}) \gamma_1^\dagger(-\vec{q}) + h.c. \}, \quad (4)$$

where $E(\pm\vec{q}) = \hbar(q_x^2 + q_z^2)/(2M)$. The γ_1 mode after the Bogoliubov diagonalization yields the spectra $\omega(\vec{q}) = \sqrt{E_{\vec{q}}(E_{\vec{q}} + 2gN_0)} \approx \sqrt{\frac{\hbar g N_0}{M}} \sqrt{q_x^2 + q_z^2}$. This is the phonon mode describing the overall density fluctuations, which exhibits linear dispersion relation for $\vec{q}$ in the xz -plane and becomes soft for $\vec{q}$ along $\hat{e}_y$. The γ_2 mode represents the relative density fluctuations between two condensates which can also be considered as spin density fluctuations. This mode remains a free particle spectrum $E(\vec{q})$. Both of the $\gamma_{1,2}$ modes, at $q \ll k_0$, only depend on the total condensation density $N_A + N_B = N_0$. Hence, the contribution from part I does not lift the degeneracy between different partitions of (N_A, N_B) to the quadratic order of q .

Now, we turn to the low energy excitations in part II where the dispersion of $\psi_{\vec{k},+}$ is degenerate with the state of $\psi_{-\vec{k},+}$ but not with $\psi_{2\vec{k}_A, B-\vec{k},+}$. The mean field Hamiltonian reads

$$H_{MF,2} = \sum_{\vec{k}} \psi_{\vec{k},+}^\dagger \psi_{\vec{k},+} \{ \epsilon(k) + \frac{g}{2}(N_0 - \Delta N \cos \phi_k) \} + g \{ \psi_{\vec{k},+}^\dagger \psi_{-\vec{k},+}^\dagger \sqrt{N_A N_B} \cos \phi_{\vec{k}} e^{i\phi_{\vec{k}}} + h.c. \}, \quad (5)$$

where $\Delta N = N_A - N_B$. The Bogoliubov spectra can be solved as $H_{MF,2} = \sum_{\vec{k}} \{ \omega(\vec{k}) (\gamma_3^\dagger(\vec{k}) \gamma_3(\vec{k}) + \frac{1}{2}) \}$ with

$$\omega(k) = \sqrt{\epsilon_{\vec{k}}(\epsilon_{\vec{k}} + gN_0) + \frac{g^2}{4}[N_0^2 \sin^2 \phi_k + (\Delta N)^2 \cos^2 \phi_k]} + \frac{g\Delta N}{2} \cos \phi_k. \quad (6)$$

After summing over $\vec{k}$ around the ring, the second term in Eq. 6 disappears. Thus the ground state zero point




FIG. 2: A) The radial density distribution of spin up and down components, and the total density distribution in the unit of N_0 at $\alpha = 4$ and $\beta = 40$ in a harmonic trap. B) The spin density distribution along the x -axis which spirals in the z - x plane at an approximate wavevector of $2k_0$, whose value at the origin is normalized to 1. The spin density in the whole plane exhibits the skyrmion configuration.

motion from Eq. 6 selects the optimal partition of $\Delta N = 0$, *i.e.*, an equal partition between two condensates.

Fragmentization of BECs is usually unstable against perturbations [26, 27]. It can be removed by non-conservation of momentum induced by spatial inhomogeneity. Now we consider a realistic system with confining trap. Our main interest is the interplay between the confining potential and the Rashba SO coupling which only lies in the xy -plane. Thus we neglect the effect of the trapping potential along the z -axis, and assume that the condensate wavefunction is uniform along this axis. We consider $V_{ex}(r) = \frac{1}{2}M\omega_T^2(x^2 + y^2)$, and define the characteristic SO energy scale for the harmonic trap as $E_{so} = \hbar\lambda_R/l$ where $l = \sqrt{\hbar/(M\omega_T)}$ is the length scale of the trap; correspondingly, we also define the dimensionless parameter $\alpha = E_{so}/(\hbar\omega_T) = lk_0$. Due to the 2D rotational symmetry, the single particle wavefunctions can be denoted by the total angular momentum $j_z = m + \frac{1}{2}$. The ground state single particle wavefunctions form a Kramer doublet as represented in polar coordinates as

$$\psi_{\frac{1}{2}}(r, \phi, z) = \begin{pmatrix} f(r) \\ g(r)e^{i\phi} \end{pmatrix}, \quad \psi_{-\frac{1}{2}}(r, \phi, z) = T\psi_{\frac{1}{2}}. \quad (7)$$

where both $f(r)$ and $g(r)$ are real functions.

Let us gain some intuition about $f(r)$ and $g(r)$ by considering the strong SO limit $\alpha \gg 1$. The Fourier components of $\psi_{\pm\frac{1}{2}}$ in momentum space mainly distribute around the low energy ring. They can be approximately

represented as

$$|\psi_{\pm\frac{1}{2}}\rangle = \int d\phi_k \int dk w(k) e^{\pm i\frac{1}{2}\phi_k} |\psi_{\pm}(\vec{k})\rangle, \quad (8)$$

where the weight $w(k)$ mainly distributes along the low energy ring with a broadening at the order of $1/l$. In other words, the harmonic potential, which reads $V_{ex} = \frac{1}{2}M\omega_T^2(i\partial_{\vec{k}} - \vec{A}(\vec{k}))^2$ in momentum representation, where $\vec{A}(\vec{k}) = i\langle\psi_{+}(\vec{k})|\partial_{\vec{k}}|\psi_{+}(\vec{k})\rangle$. For the Rashba coupling, the field strength of $\vec{A}(\vec{k})$ corresponds to a π -flux located at $\vec{k} = (0, 0)$. V_{ex} breaks the degeneracy of the plane wave states with $|\vec{k}| = k_0$ by quantizing the orbital motion along the ring as

$$\Delta E_{m+\frac{1}{2}} = \frac{1}{2}M\omega_T^2\left(\frac{m+\frac{1}{2}}{k_0}\right)^2 = \frac{1}{2\alpha^2}\hbar\omega_T\left(m+\frac{1}{2}\right)^2. \quad (9)$$

The ground state forms the Kramer doublets corresponding to $m + \frac{1}{2} = \pm\frac{1}{2}$, and the low energy azimuthal excitations are with $m + \frac{1}{2} = \pm\frac{3}{2}, \pm\frac{5}{2}, \dots$. The excitations in the radial direction of $|\vec{k}|$ change the node structures of $f(r)$ and $g(r)$, which cost the energy at the order of $\hbar\omega_T$ and thus are much higher energy states than the azimuthal excitations at $\alpha \gg 1$. For the strong SO coupling $\alpha \gg 1$, the single particle ground doublet states have nearly equal weight in the spin up and down components, *i.e.* $\int dr d\phi r |f(r)|^2 \approx \int dr d\phi r |g(r)|^2$, thus the spin moment averages to zero. When the bosons condense into one of the TR doublets, the average orbital angular momentum per particle is $\hbar/2$, *i.e.*, one spin component stays in the s -state and the other one in the p -state. This is a half-quantum vortex configuration [28–30], thus spontaneously breaking TR symmetry.

Next we turn on interactions and define the characteristic interaction energy scale $E_{int} = gN_0/(\pi l^2)$ and the dimensionless parameter $\beta = E_{int}/(\hbar\omega_T)$. We numerically solve the Gross-Pitaevskii equation

$$\left\{ -\frac{\hbar^2\nabla^2}{2M} + \hbar(-i\nabla_y\sigma_{x,\alpha\beta} + i\nabla_x\sigma_{y,\alpha\beta}) + g(\psi_\gamma^*\psi_\gamma) + \frac{1}{2}M\omega_T^2 r^2 \right\} \psi_\beta(r, \phi) = E\psi_\alpha(r, \phi) \quad (10)$$

in the harmonic trap with parameters $\alpha = 4$ and $\beta = 40$. The interaction effectively weakens the harmonic potential and does not change the orbital partial wave structure of the wavefunctions. We show the radial density profiles of both spin components $|f(r)|^2$ and $|g(r)|^2$ in Fig. 2 A. Each of them exhibits oscillations at a pitch value of approximately $2k_0$, which originate from the low energy ring structure and, thus, are analogous to the Friedel oscillations in fermion systems.

The spin density, defined as $\vec{S}(r, \phi) = \psi_\alpha^*(r, \phi)\vec{\sigma}_{\alpha\beta}\psi_\beta(r, \phi)$, exhibits an interesting topological spin texture configuration. Let us first look at its distribution along the x -axis where the supercurrent is

along the y -direction and the spin lies in the xz -plane. Explicitly, we express $S_z(r, \phi) = \frac{1}{2}(|f(r)|^2 - |g(r)|^2)$ and $S_x(r, \phi) = f(r)g(r)$. The radial oscillations of $|f(r)|^2$ and $|g(r)|^2$ have an approximate π phase shift, which arises from the different angular symmetries. As a result, $\vec{S}$ spirals along the x -axis as plotted in Fig. 2 B at the pitch value of the density oscillations. The spin density distribution in the whole space can be obtained through a rotation around the z -axis, which exhibits the skyrmion configuration.

The recent research focus of the “synthetic gauge fields” in cold atom systems provides a promising method to observe the above proposed exotic spontaneous half-quantum vortex [10–12, 14]. Artificial magnetic fields have been synthesized by light-atom interaction to generate vortices experimentally [12]. Similar methods to generate SO coupling have been proposed [10, 13, 31], and are currently under experimental implementation. A convenient method is the “tripod scheme”: three atomic degenerate hyperfine ground states are coupled to an excited state by laser fields to create a pair of degenerate dark states denoted with pseudospin indices $|\uparrow\rangle$ and $|\downarrow\rangle$. The spatially dependent laser fields couple the pseudospin and the momentum of the atoms. A typical set of parameter values are provided in Ref. [10] for a trap frequency $\omega_T = 2\pi \times 10$ Hz, a characteristic Rabi frequency $\Omega = 2\pi \times 10^7$ Hz, and a detuning $\Delta = 2\pi \times 10^{11}$ Hz, which satisfy $\omega_T \ll \Omega \ll \Delta$ and $\omega_T \ll \Omega^2/\Delta$. The value of α can vary from 0 up to the order of several tens. With the typical particle number $N_0 = 10^6$, the interaction energy scale E_{int} is around $100nK$, or $2kHZ$ [32], and thus β is of the order of several tens. Hence, the parameters used in the Fig. 2 A and B are realistic in real experiments. The experimental detection would be straightforward as performed in previous time-of-flight imaging in vortex experiments [33, 34]. By separately imaging the density profiles of the pseudospin up and down components, the radial oscillation of the S_z component can be directly seen.

Another class of boson systems exhibiting SO coupling is the indirect excitons in 2D coupled double quantum well systems. We only need to consider the heavy hole (hh) band with the effective mass m_{hh}^* and $j_z = \pm\frac{3}{2}\hbar$, which is separated from the light hole band with a gap of the order of $10meV$. The Hamiltonians of the electron-hole system reads $H_e = -\frac{\hbar^2}{2m_e^*}\nabla_e^2 + i\hbar\lambda_R(\partial_{e,x}\sigma_y - \partial_{e,y}\sigma_x)$, $H_{hh} = -\frac{\hbar^2}{2m_{hh}^*}\nabla_h^2$, and $H_{e-ph} = -\frac{e^2}{\varepsilon\sqrt{|\vec{r}_e - \vec{r}_{hh}|^2 + d^2}}$, where m_e^* is the effective mass of the conduction electrons; λ_R is the Rashba SO coupling strength of the conduction electron; ε is the dielectric constant; and d is the distance between the double wells. The center-of-mass motion of excitons in the BEC limit can be separated from the electron-hole relative motion. Similarly to Ref. [35], the effective Hamiltonian of the 4-component hh excitons denoted as $(s_e, j_{hh}) = (\pm\frac{1}{2}, \frac{3}{2})$, $(\pm\frac{1}{2}, -\frac{3}{2})$ can be represented

by the matrix form as

$$H_{ex} = \begin{pmatrix} E_{ex}(\vec{k}) & H_{so}(\vec{k}) & 0 & 0 \\ H_{so}^*(\vec{k}) & E_{ex}(\vec{k}) + \Delta(\vec{k}) & W(\vec{k}) & 0 \\ 0 & W^*(\vec{k}) & E_{ex}(\vec{k}) + \Delta(\vec{k}) & H_{so}(\vec{k}) \\ 0 & 0 & H_{so}^*(\vec{k}) & E_{ex}(\vec{k}) \end{pmatrix},$$

where $\vec{k}$ is the center-of-mass momentum; $M = m_e^* + m_{hh}^*$ is the total mass of the exciton; $E_{ex}(\vec{k}) = \hbar^2 k^2 / (2M)$; $H_{so}(\vec{k}) = -\frac{m_e^*}{M}\lambda_R(k_y + ik_x)$; $\Delta(\vec{k})$ is the exchange integral and $W(\vec{k})$ has the d -wave structure as $(k_x + ik_y)^2$, both of which are exponentially suppressed by the tunneling barrier between electron and hole layers, and can be neglected. Consequently, H_{ex} becomes block-diagonalized. We consider using circularly polarized light to pump the exciton of $(-\frac{1}{2}, \frac{3}{2})$, and then focus on the left-up block with heavy hole spin $j_z = \frac{3}{2}$. We will then end up with the 2D version of the Hamiltonian given by Eq. 1.

In summary, we find that bosons with SO coupling exhibit a novel type of BEC beyond Feynman’s argument. In the limit of large SO coupling, the degeneracy of the single particle ground states in momentum space leads to two fragmented condensations with equal particle number partition in homogeneous space. In the presence of a harmonic trap, the condensate spontaneously breaks TR symmetry and develops the half-quantum vortex configuration with the skyrmion type topological spin density configurations.

C. W. is supported by NSF under No. DMR- 0804775, and ARO-W911NF0810291.

-
- [1] R. P. Feynman, *Statistical Mechanics, A Set of Lectures* (Addison-Wesley Publishing Company, ADDRESS, 1972).
 - [2] T. Mueller, S. Foelling, A. Widera, and I. Bloch, *Phys. Rev. Lett.* **99**, 200405 (2007).
 - [3] A. Isacsson and S. M. Girvin, *Phys. Rev. A* **72**, 053604 (2005).
 - [4] W. V. Liu and C. Wu, *Phys. Rev. A* **74**, 13607 (2006).
 - [5] C. Wu, W. V. Liu, J. E. Moore, and S. Das Sarma, *Phys. Rev. Lett.* **97**, 190406 (2006).
 - [6] A. B. Kuklov, *Phys. Rev. Lett.* **97**, 110405 (2006).
 - [7] V. M. Stojanovic, C. Wu, W. V. Liu, and S. D. Sarma, *Phys. Rev. Lett.* **101**, 125301 (2008).
 - [8] G. Juzeliunas *et al.*, *Phys. Rev. Lett.* **100**, 200405 (2008).
 - [9] J. Y. Vaishnav and C. W. Clark, *Phys. Rev. Lett.* **100**, 153002 (2008).
 - [10] T. D. Stanescu, C. Zhang, and V. Galitski, *Phys. Rev. Lett.* **99**, 110403 (2007).
 - [11] Y.-J. Lin *et al.*, *Phys. Rev. Lett.* **102**, 130401 (2009).
 - [12] Y.-J. Lin *et al.*, *Nature* **462**, 628 (2009).
 - [13] I. B. Spielman, *Phys. Rev. A* **79**, 063613 (2009).
 - [14] T. Stanescu, B. Anderson, and V. Galitski, *Phys. Rev. A* **78**, 023616 (2008).
 - [15] D. W. Snoke, J. P. Wolfe, and A. Mysyrowicz, *Phys. Rev. B* **41**, 11171 (1990).

- [16] L. V. Butov, J. Phys.: Cond. Matt. **16**, R1577 (2004).
- [17] L. V. Butov, J. Phys.: Cond. Matt. **19**, 295202 (2007).
- [18] L. A. V. Timofeev V. B., Gorbunov A. V., J. Phys.: Cond. Matt. **19**, 295209 (2007).
- [19] T. Hakioglu and M. Sahin, Phys. Rev. Lett. **98**, 166405 (2007).
- [20] M. A. Can and T. Hakioglu, arXiv:0808.2900, 2008.
- [21] W. Yao and Q. Niu, Phys. Rev. Lett. **101**, 106401 (2008).
- [22] L. V. Butov *et al.*, Phys. Rev. Lett. **73**, 304 (1994).
- [23] L. V. Butov and A. I. Filin, Phys. Rev. B **58**, 1980 (1998).
- [24] L. V. Butov *et al.*, Phys. Rev. Lett. **86**, 5608 (2001).
- [25] L. V. Butov, A. C. Gossard, and D. S. Chemla, Nature **418**, 751 (2002).
- [26] E. J. Mueller, T. L. Ho, M. Ueda, and G. Baym, Phys. Rev. A **74**, 33612 (2006).
- [27] T. L. Ho and S. K. Yip, Phys. Rev. Lett. **84**, 4031 (2000).
- [28] M. M. Salomaa and G. E. Volovik, Phys. Rev. Lett. **55**, 1184 (1985).
- [29] C. Wu, J. P. Hu, and S. C. Zhang, Int. J. Mod. Phys. B **V24**, 311 (2010).
- [30] F. Zhou, Int. Jour. Mod. Phys.B, **17** **17**, 2643 (2003).
- [31] J. Larson and E. Sjöqvist, Phys. Rev. A **79**, 043627 (2009).
- [32] A. J. Leggett, Rev. Mod. Phys. **73**, 307 (2001).
- [33] K. W. Madison, F. Chevy, W. Wohlleben, and J. Dalibard, Phys. Rev. Lett. **84**, 806 (2000).
- [34] B. P. Anderson, P. C. Haljan, C. E. Wieman, and E. A. Cornell, Phys. Rev. Lett. **85**, 2857 (2000).
- [35] M. Z. Maialle, E. A. de Andrada e Silva, and L. J. Sham, Phys. Rev. B **47**, 15776 (1993).